

Pain Pre-Procedure Instructions

Please read the following instructions carefully to ensure that you can complete your scheduled procedure and avoid cancellation/rescheduling.

BEFORE THE PROCEDURE:

- Please notify Texas Regional Clinic if you are **allergic** to iodine, seafood, shellfish, latex, steroid, local anesthetics (i.e. lidocaine), or any medication in general.
- If you have diabetes, please **continue** your recommended diet and **continue** to take your diabetic medication as prescribed by your doctor.
- If you have high blood pressure, please continue to take your blood pressure medication as prescribed by your doctor.
- Nothing to eat or drink (shake, smoothie, soup, coffee, etc) at least 8 hours prior to the procedure. You can drink water up until 2 hours before the procedure.
- > Schedule a driver to be present at the time of check-in on the day of the procedure.
 - If you plan to use Lyft or Uber, please have another adult accompany you on the ride.
 - If you do not have an adult with you, then you <u>CANNOT</u> receive any form of sedation.
- ➤ If you're taking blood thinning (anti-coagulation/anti-platelet) medication prescribed by a physician (i.e. Plavix, Aspirin 325 mg, Warfarin, Eliquis, Xarelto, Lovenox, Heparin, etc.) We will need to obtain a **clearance** from your prescriber **before stopping your medication**.
- ➤ Do **NOT** take these over-the-counter medications for at least <u>7 days prior</u> to the procedure and 24-48 hours after: Alka-Seltzer, BC Powder, Excedrin, or anything with Aspirin.
- ➤ If you are taking Baby Aspirin(81mg) for preventative, please stop <u>5 days prior</u> to the procedure and do not resume until <u>24-48 hours after</u>.
- For procedures involving the spine, please Do NOT take anti-inflammatory medications, such as Advil/Motrin/Ibuprofen, Aleve/Naproxen, Mobic/Meloxicam, Diclofenac, etc., for at least 48 hours prior to the procedure. Your provider may require you to stop for longer than 48 hours, depending on your risk factors. (Over-the-counter Tylenol is okay to take).
- ➤ Do **NOT** take any type of narcotic pain medications, benzodiazepines, or muscle relaxers on the day of the procedure.
- Please notify us if you have any active infection or if you are taking any antibiotics. We will have to schedule the procedure at least a week after you have completed your antibiotics.
- ➤ If you have recently received a COVID-19 vaccine/booster, we strongly recommend you wait 2 weeks before proceeding with a steroid injection.
- ▶ If you are planning on receiving a COVID-19 vaccine/booster, we strongly recommend you wait 2 weeks after you have received a steroid injection.

If you have any question or concern, please call **Texas Regional Clinic** at 713-498-1741.



Pain Pre-Procedure Questionaire

Please answer the following questions to the best of your knowledge so that we can help you schedule your procedures as soon as possible.

1. Are you currently taking any blood thinning (anti-coagulation/anti-platelet) medication prescribed by a physician (i.e. Plavix, Aspirin, Warfarin, Eliquis, Xarelto, Lovenox, Heparin, etc.)	NO	YES
If yes, please fill out the following:		
Medication name: Dosage:		
Prescriber/Clinic name:		
Clinic Address:		
Clinic Telephone #:		
Clinic fax #:		
2. Are you taking Alka-Seltzer, BC Powder, Excedrin, or anything with Aspirin?	NO	YES
3. Are you taking any anti-inflammatory medication? (i.e. Advil/Motrin/Ibuprofen, Aleve/Naproxen, Mobic/Meloxicam, Diclofenac, etc.)	NO	YES
4. Are you allergic to iodine, seafood, shellfish, latex, steroid, or local anesthetic (i.e. lidocaine, mepivacaine, bupivacaine) ? If yes, which of the above:	NO	YES
5. Do you currently have an active infection or are undergoing treatment with antibiotics for an infection?	NO	YES
6. Are you currently taking oral steroid?	NO	YES

Note to staff: Please fax this form with the referral.